



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

76; VIII: Germany, 71-129; IX: Hungary, 130-137; X: Italy, 138-143; XI: Japan, 144-152; XII: The Netherlands, 153-157; XIII: Rumania, 158-159; XIV: Russia, 160-167; XV: Spain, 168-170; XVI: Sweden, 171-190; XVII: Switzerland, 191-199; XVIII: The United States, 200-211; XIX: Summary and comparative remarks, 212-230. Appendices, A, B, England: Cambridge local examinations senior students; Oxford and Cambridge schools; examination board; University of London matriculation examinations; Entrance scholarship examination papers, Cambridge, 231-251; C, D, France: Concours for admission to the Ecole Normale Supérieure, and for the bourses de licence in 1913, Agrégation des sciences mathématiques, 252-266; E, Germany: Reifeprüfungen, Lehramtsprüfungen, 267-274; F, Japan: Examination questions, 275-278. Index, 279-289.

## QUESTIONS AND DISCUSSIONS.

Edited by U. G. MITCHELL, University of Kansas, Lawrence.

Three questions which, among others, have been standing for some time without having been answered are republished below in the hope that some of our readers may thereby be stimulated to send in suitable replies.

**15.** In the *Proceedings of the Royal Society of Edinburgh*, Vol. VII, p. 144, in some mathematical notes by Professor P. G. Tait, it is stated:

"If  $x^3 + y^3 = z^3$ , then  $(x^3 + z^3)^3 y^3 + (x^3 - y^3)^3 z^3 = (z^3 + y^3)^3 x^3$ .

"This furnishes an easy proof of the impossibility of finding two integers the sum of whose cubes is a cube."

How does this "easy proof" follow?

(A partial reply to the above has been received showing that if  $x^3 + y^3 = z^3$ , then  $(x^3 + z^3)^3 y^3 + (x^3 - y^3)^3 z^3 = (z^3 + y^3)^3 x^3$ . Can some one show how the "easy proof" then follows?)

**21.** For the diophantine equation

$$x^2 - y^3 = 17$$

there are known the following solutions:

$$x = 3, \quad 4, \quad 5, \quad 9, \quad 23, \quad 282, \quad 375, \quad 378661,$$

$$y = -2, \quad -1, \quad 2, \quad 4, \quad 8, \quad 43, \quad 52, \quad 5234.$$

One of our readers, who supplied the foregoing facts, desires to know the answers to the following questions: Are there other solutions of the given diophantine equation? How may all the solutions of this equation be found by a systematic procedure?

**32.** In a discussion of the Peaucellier<sup>1</sup> Cell by analytic methods the following equations are obtained:

$$(1) \quad (x_2 - x_1)^2 + (y_2 - y_1)^2 - b^2 = 0; \quad (2) \quad (x_3 - x_1)^2 + (y_3 - y_1)^2 - b^2 = 0;$$

$$(3) \quad (x_2 - X)^2 + (y_2 - Y)^2 - b^2 = 0; \quad (4) \quad (x_3 - X)^2 + (y_3 - Y)^2 - b^2 = 0;$$

$$(5) \quad x_2^2 + y_2^2 - K^2 = 0; \quad (6) \quad x_3^2 + y_3^2 - K^2 = 0;$$

$$(7) \quad x_1^2 + y_1^2 - 2cx_1 = 0.$$

The result of eliminating  $x_1, y_1, x_2, y_2, x_3, y_3$  gives an equation of the first degree, which establishes that the linkage will trace a straight line. There are various ways of effecting this elimination.

1. What element of the situation is left unused by the following procedure in the elimination?

(a) From equations (1), (3), (5) eliminate  $x_2$  and  $y_2$  and obtain an equation

$$(8) \quad f_1(x_1, y_1) = 0.$$

(b) From equations (2), (4), (6) eliminate  $x_3$  and  $y_3$  and obtain an equation

$$(9) \quad f_2(x_1, y_1) = 0.$$

(c) From equations (7), (8), (9) eliminate  $x_1$  and  $y_1$  and obtain the desired equation.

2. How should this procedure be supplemented to secure the result?

<sup>1</sup> If reference is made to the article on "Linkages" in the December, 1915, MONTHLY, by Mr. Leavens, the following coordinates may be applied to his figure:  $O(0, 0)$ ;  $C(c, 0)$ ;  $P_1(x_1, y_1)$ ;  $M(x_2, y_2)$ ;  $M_1(x_3, y_3)$ ;  $P_2(X, Y)$ .